



Original Investigation | Psychiatry

Risk and Protective Factors for Suicide Mortality in Youths

A Systematic Review and Meta-Analysis

Marie-Claude Geoffroy, PhD; Elizabeth Baker-Sullivan, MScPH; Nusrat Jahan Nitu, MBBS; Alessia Civita, BA; Jill Boruff, MLIS, AHIP; Mark Sinyor, MSc, MD; Jo Robinson, PhD; Kathleen MacDonald, PhD; Lauren Anzarouth, MSc; Elise Chartrand, MSc; Srividya N. Iyer, PhD; Rachel Langevin, PhD; Jane Pirkis, PhD; Brett Thombs, PhD

Abstract

IMPORTANCE Risk and protective factors for suicide mortality in youths remain poorly synthesized, as prior reviews have focused on all ages or nonfatal outcomes.

OBJECTIVE To systematically assess factors associated with risk of suicide mortality in youths.

DATA SOURCES MEDLINE, PsycINFO, Embase, and CINAHL from inception to March 7, 2025.

STUDY SELECTION Case-control and cohort studies of youths (aged ≤ 24 years) examining risk and/or protective factors associated with suicide mortality vs living general-population controls were included. Two independent reviewers screened 9497 records.

DATA EXTRACTION AND SYNTHESIS Following Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines, 2 reviewers independently screened reports; 1 extracted data, verified by a second. Evidence was synthesized using vote counting and random-effects meta-analysis in April 2026.

MAIN OUTCOMES AND MEASURES The primary outcome was suicide mortality at age 24 years or younger.

RESULTS Ninety reports from 68 studies, mostly from high-income countries, identified distinct risk and/or protective factors; 54 reports contributed to 30 meta-analyses. The factors associated with the highest odds of suicide risk included schizophrenia (odds ratio [OR], 22.23; 95% CI, 12.05-41.03; $I^2 = 85.4\%$; 7 reports), mood disorders (OR, 11.32; 95% CI, 6.11-20.97; $I^2 = 64.6\%$; 7 reports), and self-harm (OR, 14.06; 95% CI, 5.58-35.39; $I^2 = 90.1\%$; 10 reports). Clinical indicators of health care use were also associated with higher risk, including mental health services use in preceding year (OR, 7.39; 95% CI, 6.45-8.47; $I^2 = 0.0\%$; 5 reports) and psychiatric admission (OR, 31.96; 95% CI, 13.83-73.86; $I^2 = 94.8\%$; 6 reports). At the socioecological level, several indicators were associated with higher risk, including maltreatment (OR, 4.03; 95% CI, 1.41-11.50; $I^2 = 67.1\%$; 5 reports), out-of-home placement (OR, 4.47; 95% CI, 2.15-9.28; $I^2 = 42.1\%$; 5 reports), youth justice system involvement (OR, 2.70; 95% CI, 1.94-3.75; $I^2 = 64.2\%$; 7 reports), and low educational attainment (OR, 2.95; 95% CI, 1.66-5.24; $I^2 = 76.0\%$; 5 reports). In contrast, indicators of family stability were associated with lower risk, including living with both parents (OR, 0.55; 95% CI, 0.48-0.62; $I^2 = 12.1\%$; 11 reports). Heterogeneity was substantial across analyses, while Newcastle-Ottawa ratings indicated moderate-to-high study quality.

(continued)

Key Points

Question What factors are associated with risk of suicide mortality among individuals aged 24 years and younger?

Findings In this systematic review and meta-analysis of 90 reports (68 reports), mental disorders—particularly acute conditions such as schizophrenia and mood disorders—and prior self-harm were associated with suicide mortality. Prior health care contact (eg, psychiatric admission) and a range of social determinants (eg, maltreatment, out-of-home placement, and social disadvantage) were also associated with higher risk, whereas family stability (eg, living with both parents) was associated with lower risk.

Meaning Suicide mortality in youths reflects both psychiatric vulnerability and social adversity, supporting the need for integrated prevention strategies combining early intervention with population-level prevention approaches targeting social determinants.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY-NC-ND License, which does not permit alteration or commercial use, including those for text and data mining, AI training, and similar technologies.

Abstract (continued)

CONCLUSIONS AND RELEVANCE In this systematic review and meta-analysis, suicide mortality in youths was associated with mental disorders, health care contact, and adversity, supporting both clinical care and population-level prevention, with future research needed in underrepresented populations.

JAMA Network Open. 2026;9(7):e2621622. doi:10.1001/jamanetworkopen.2026.21622

Introduction

Suicide remains one of the leading causes of death among young people aged 10 to 24 years worldwide. Recent World Health Organization mortality data from 52 countries indicate an upward trend in many high-income countries—especially among girls—whereas boys continue to die by suicide at 2 to 5 times the rate of girls.¹ Yet the risk and protective factors for suicide mortality remain insufficiently documented in youths, limiting our ability to identify developmentally specific targets for prevention.

Youth-focused reviews have largely examined suicidal ideation and self-harm rather than suicide mortality, typically focusing on a limited set of factors,^{2,3} whereas only a few capture a broader range of risk and protective factors.^{4,5} Given that suicidal ideation and self-harm are far more prevalent than suicide death, findings from these reviews may not fully inform suicide mortality prevention efforts. Conversely, reviews of suicide mortality typically pool evidence across all ages,⁶⁻⁸ overlooking key developmental differences in youth.⁹ One prior review examined sex-stratified risk and protective factors for suicidal behaviors in youths and included suicide death, but was based on a very limited number of studies.¹⁰

Youth suicide mortality may reflect the convergence of mental disorders—captured through psychiatric diagnoses or health care contact—and accumulated social adversity, such as abuse and poverty. However, to our knowledge, no prior systematic review and meta-analysis has synthesized the breadth of risk and/or protective factors for suicide mortality among young people aged 24 years or younger. To directly inform prevention strategies targeting youth suicide deaths, we systematically reviewed and meta-analyzed risk and protective factors for suicide mortality among individuals aged 24 years or younger.

Methods

In this systematic review and meta-analysis, we included primary studies that (1) employed case-control studies (psychological autopsy or registered-based studies) or cohort (retrospective or prospective) studies, (2) quantitatively assessed risk and/or protective factors by comparing youths aged 24 years or younger who died by suicide with living controls, and (3) reported suicide mortality separate from other outcomes such as nonfatal self-harm. The review was preregistered with PROSPERO (CRD420251008539) and reported in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) reporting guidelines. We deviated from the initial protocol by not applying a lower age restriction of 10 years for suicide, because age-specific information was not consistently reported across studies. Studies with broad age ranges were included if at least 80% of participants were aged younger than 25 years and the full sample fell within a 0- to 30-year range, or if the mean age was younger than 25 years.

A health sciences librarian (J.B.) searched MEDLINE (Ovid), PsycINFO (Ovid), Embase (Ovid), and CINAHL (EBSCO) for articles published from inception to March 7, 2025. The search combined terms related to suicide mortality, child, adolescent, and/or young adult and case-control or cohort studies with no restrictions applied to language or publication date. It was peer-reviewed by a second librarian (not a coauthor of this study) using the Peer Review of Electronic Search Strategies; see Boruff¹¹ for the search strategy.

Titles and abstracts were uploaded into Covidence; duplicates were removed. Two independent reviewers (M.C.G or E.B.S.; and N.J.N, A.C., K.M., or L.A.) evaluated titles and abstracts and then full text review. Any discrepancies were resolved through consensus, with a third investigator (M.C.G or E.B.S.) consulted as necessary. Standardized screening guides (eAppendixes 1 and 2 in [Supplement 1](#)) were developed and refined through an iterative process to ensure consistency and accuracy. All team members received training across multiple sessions. Because articles in all languages were eligible, non-English records were assessed using Google Translate or fluent speakers. As this study is a systematic review of previously published literature, no new ethical approval was required.

Data Extraction

Data were extracted from the full text and supplementary materials. Extracted information included first author, publication year, country, number of suicide deaths and controls (or total living cohort), mean age and age range at the time of suicide death, year(s) in which suicide deaths occurred, the proportion of suicide deaths among male youths, and all risk and protective factors, with no a priori selection of factors. We used World Bank classifications for country income and region. Procedures used to handle duplicate and/or overlapping risk and protective factors across reports are described in the eMethods in [Supplement 1](#). Study quality was assessed using the Newcastle-Ottawa Scale, which evaluates studies across 3 dimensions: selection of study groups, comparability of groups, and ascertainment of exposure or outcome,¹² for both cohort and case-control studies (eAppendixes 3 and 4 in [Supplement 1](#)). Data extracted for synthesis are described below. For all extractions, 1 reviewer (M.C.G or E.B.S.) extracted data using a standardized form, and a second reviewer (A.C., E.C., or N.J.N.) independently verified their accuracy and completeness.

Statistical Analysis


First, all risk and protective factors were grouped into 5 broad a priori domains: mental health and self-harm, physical disorders, health care service contacts, prenatal and perinatal factors, and socioecological determinants (classified using the American Psychiatric Association Social Determinants of Mental Health framework).¹³ Second, within each domain, factors examined in at least 2 reports of 2 different studies (eg, completed or current level of education and highest grade completed) were grouped into comparable risk and protective factors for narrative synthesis (eTables 1-5 in [Supplement 1](#)). When a report included more than 1 related measure for the same risk and protective factors (eg, continuous score and yes vs no), we retained the measure that was most similar to those used in other reports (eMethods in [Supplement 1](#)). Third, for factors examined in at least 2 reports, we used vote counting based on the direction of association in line with Cochrane Handbook guidance.¹⁴ This narrative synthesis approach summarizes evidence by counting studies according to whether associations were harmful, protective, or null, based on direction, with factors for which direction could not be determined (ie, no estimates reported) coded as not reported. Fourth, for factors examined in at least 5 reports and that could be harmonized (eTables 6-15 in [Supplement 1](#)), we performed random-effects meta-analyses using restricted maximum likelihood estimation in the meta package in R statistical software version 4.1.2 (R Project for Statistical Computing) in April 2026. For both vote-counting and meta-analysis, we prioritized unadjusted or minimally adjusted estimates for sex and age. Estimates extracted included odds ratios (ORs), hazard ratios (HRs), and risk ratios (RRs), or ORs were calculated from 2 × 2 tables, with corresponding 95% CIs (eMethods in [Supplement 1](#)). ORs were retained for meta-analyses, with RRs and HRs approximated to ORs.^{15,16} Statistical significance was defined as a 95% CI that did not include the null value of 1. Forest plots were used to represent results graphically. We assessed heterogeneity using the I^2 statistic and 95% prediction intervals, calculated from the random-effects model and representing the expected range of effect estimates in a new study. Potential publication bias was evaluated using funnel plots. Sensitivity analyses included stratification by study design (case-control vs cohort).

Results

We identified 14 238 records through database searches, screened 9497 after removing duplicates, and reviewed 1573 full-text documents, with an additional 34 retrieved from the reference lists of secondary sources (eg, reviews and commentaries). In total, 154 reports from 120 studies met our inclusion criteria. We further excluded 56 reports due to limited generalizability of selected populations (eg, youth involved in child welfare or in contact with health care services) to enhance generalizability to the broader population, and 6 reports on pharmacological treatments, as these were beyond the scope of this review and are summarized elsewhere, and 2 duplicate reports (eAppendix 5 in [Supplement 1](#) for full list of excluded articles after full-text screening). Ultimately, we included 90 reports¹⁷⁻¹⁰⁶ from 68 studies in the systematic review (**Figure 1**).

The 68 included studies were conducted primarily in high-income countries (78 reports for 58 studies^{17-33,38-62,65-67,69,73,75-105}) across 3 World Bank regions (North America, Europe and Central Asia, and East Asia and Pacific) with most from Sweden (17 reports of 15 studies^{19,23,24,39,45,47,48,62,66,75,87,96,98,101,102,104,105}) and the US (27 reports of 13 studies^{17,18,26-31,42,49-53,56-58,60,67,73,77,79,90-94}). An additional 10 studies were from upper-income and lower-middle-income countries, mainly China (including Taiwan and Hong Kong, 11 reports of 9 studies^{34-37,63,64,70-72,74,106}) and Sri Lanka (1 study⁶⁸). We identified 36 case-control studies (27 reports of 12 psychological autopsy studies^{19,22,26-31,34,42,55-58,61,70,83-85,89-94,102,106} and 26 reports of 24 registered-based studies^{18,20,21,32,35,36,38,41,43,44,60,63,64,67,72-74,76,77,79-82,86,95,100}) and 37 reports of 32 cohort studies^{17,23-25,33,37,39,40,45-54,59,62,65,66,68,69,71,75,78,87,88,96-99,101,103-105} (mostly retrospective). Among these, 16 reports of 7 studies^{18,26-31,42,57,58,61,77,90-93} were published before 2000. The number of suicides ranged from 4 to 4828, with only 2 studies^{17,18} including fewer than 10 suicides. Study characteristics are summarized in eTable 16 in [Supplement 1](#).

Figure 1. Flowchart of the Identification and Selection of Studies on Risk and/or Protective Factors for Youth Suicide Mortality



Narrative Synthesis

Of 846 extracted factors, 106 distinct factors examined in at least 2 reports were included in vote counting: 25 mental health disorders and self-harm, 8 physical health disorders, 8 health care service contact factors, 6 prenatal and perinatal factors, and 59 socioecological determinants spanning socioeconomic opportunities for youths, families, and communities, the physical environment, highly detrimental societal problems, stigma and discrimination, stresses and traumas associated with immigration, and protective factors in accordance with American Psychiatric Association framework. Factors included and excluded from vote counting are reported per domain in eTables 1 to 5 in [Supplement 1](#). The most frequently studied risk factors (each examined in more than 10 studies) were depression, self-harm, substance abuse, and any mental health disorders. Across factors, most associations were in a harmful direction, whereas only 2 were clearly protective within the social determinant's domain—living with both parents and having married parents (see [Figure 2](#) for social determinants, and eFigure 1 and eTable 17 in [Supplement 1](#) for other factors).

Meta-Analyses

Overall, 54 reports^{17-20, 22-27, 29, 31, 34-42, 49, 50, 55, 58-61, 64, 66, 67, 72, 75-81, 83, 84, 86, 89, 91-95, 97, 99-101, 105, 106} from 45 unique studies were included in the meta-analysis, comprising 29 case-control studies (including 19 reports of 12 psychological autopsy studies^{19, 22, 26, 27, 29, 31, 34, 42, 55, 58, 61, 83, 84, 89, 91-94, 106} and 18 reports of 17 register-based studies^{18, 20, 35, 36, 38, 41, 60, 64, 67, 72, 76, 77, 79-81, 86, 95, 100}) and 17 reports of 16 cohort studies^{17, 23-25, 37, 39, 40, 49, 50, 59, 66, 75, 78, 97, 99, 101, 105} from 13 countries (0 in lower-middle-income countries). Together, they examined 30 distinct risk factors and protective factors across 4 broad domains: mental disorders, physical disorders, health care contacts, and socioecological determinants. Study quality was rated as moderate in 7 reports of 5 case-control studies,^{18, 42, 55, 67, 83, 84, 89} high in 30 reports of 24 case-control studies,^{19, 20, 22, 26, 27, 29, 31, 34-36, 38, 41, 58, 60, 61, 64, 72, 76, 77, 79-81, 86, 91-95, 100, 106} and high in 17 reports of 16 cohort studies.^{17, 23-25, 37, 39, 40, 49, 50, 59, 66, 75, 78, 97, 99, 101, 105} Reports included in the meta-analysis are in bold in eTable 16 in [Supplement 1](#), and risk of bias assessments are summarized in eTables 18 and 19 in [Supplement 1](#). Funnel plots were visually inspected for asymmetry (eFigure 2 in [Supplement 1](#)). However, given the small number of reports for most meta-analyses (<10), funnel plot interpretation was limited.¹⁰⁷ Sensitivity analyses stratified by study design (case-control vs cohort) showed no clear differences between designs (eTable 20 in [Supplement 1](#)).

Mental Disorders and Self-Harm

Among mental disorder domains, 11 disorders were meta-analyzed (≥ 5 reports each). Associations with youth suicide mortality existed across nearly all disorders, with particularly elevated odds for schizophrenia (OR, 22.23; 95% CI, 12.05-41.03; $I^2 = 85.4\%$; 7 reports),^{26, 37, 64, 72, 80, 89, 91} bipolar disorder (OR, 12.40; 95% CI, 3.17-48.47; $I^2 = 75.4\%$; 5 reports),^{26, 64, 72, 83, 91} any unspecified mood disorders (OR, 11.32; 95% CI, 6.11-20.97; $I^2 = 64.6\%$; 7 reports),^{22, 26, 34, 61, 89, 91, 93} and depression (OR, 8.34; 95% CI, 3.44-20.19; $I^2 = 94.3\%$; 10 reports),^{18, 19, 26, 41, 64, 72, 76, 83, 91, 93} but heterogeneity was high. Borderline personality disorder was examined in only 1 psychological autopsy study in youths aged 18 to 25 years (OR, 12.66; 95% CI, 4.46-35.94).¹⁹ Alcohol abuse (OR, 5.13; 95% CI, 4.44-5.93; $I^2 = 0.0\%$; 6 reports)^{19, 26, 67, 83, 91, 94} and drug use (OR, 6.35; 95% CI, 2.98-13.52; $I^2 = 0.0\%$; 5 reports)^{19, 26, 49, 83, 91} were also associated with suicide mortality, with no observed heterogeneity, in addition to substance use (OR, 7.21; 95% CI, 3.85-13.50; $I^2 = 90.2\%$; 9 reports).^{19, 26, 34, 38, 61, 64, 83, 91, 93} Associations were observed for all other disorders. Prior self-harm was also associated with suicide mortality (OR, 14.06; 95% CI, 5.58-35.39; $I^2 = 90.1\%$; 10 reports)^{22, 23, 31, 41, 50, 61, 84, 91, 92, 94} ([Figure 3](#); eTable 21 in [Supplement 1](#)).

Figure 2. Bar Graphs of Number of Reports Reporting Harmful, Protective, Null, or Unreported Associations Between Social Determinants and Youth Suicide Mortality (Vote-Counting Synthesis)



Factors are grouped by domain. See eFigure 1 in Supplement 1 for other risk and protective factors.

Physical Disorders

Across 5 reports,^{34,38,72,93,106} any physical disorder was associated with higher suicide mortality risk (OR, 1.46; 95% CI, 0.68-3.12). However, the estimate was imprecise and highly heterogeneous ($I^2 = 67.2\%$) (eTable 22 in Supplement 1).


Health Care Service Contacts

Among health care service contacts, 3 exposures were meta-analyzed. Associations with youth suicide mortality were observed for any inpatient psychiatric admission (OR, 31.96; 95% CI, 13.83-73.86; $I^2 = 94.8\%$; 6 reports).^{19,20,26,72,86,94} Any contact with mental health care was also associated with increased odds of suicide mortality, particularly when occurring within the year preceding death (OR, 7.39; 95% CI, 6.45-8.47; $I^2 = 0.0\%$; 5 reports)^{26,84,86,91,94} compared with lifetime exposure (OR, 3.56; 95% CI, 2.71-4.67; $I^2 = 0.1\%$; 9 reports)^{19,22,26,50,55,84,91,92,94} (Figure 4 and eTable 23 in Supplement 1).

Socioecological Determinants


Among the 59 socioecological determinants, 14 were meta-analyzed. Most focused on socioeconomic adversity. The highest odds of youth suicide mortality were observed for maltreatment (OR, 4.03; 95% CI, 1.41-11.50; $I^2 = 67.1\%$; 5 reports)^{19,31,38,58,92} and out-of-home-placement (OR, 4.47; 95% CI, 2.15-9.28; $I^2 = 42.1\%$; 5 reports).^{19,40,49,61,79} Justice system involvement for youths (OR, 2.70; 95% CI, 1.94-3.75; $I^2 = 64.2\%$; 7 reports)^{17,19,27,50,58,60,100} and parents (OR, 2.53; 95% CI, 1.38-4.64; $I^2 = 60.4\%$; 5 reports)^{19,24,29,40,58} were also associated with

Figure 3. Forest Plot of Pooled Associations of Mental Disorders and Prior Self-Harm With Youth Suicide Mortality (Meta-Analysis)



ADHD indicates attention-deficit/hyperactivity disorder; OR, odds ratio.

Figure 4. Forest Plot of Pooled Associations of Health Care Contact With Youth Suicide Mortality (Meta-Analysis)



OR indicates odds ratio.

higher risk. Indicators of family stability were consistently associated with lower risk, including living with both parents (OR, 0.55; 95% CI, 0.48-0.62; $I^2 = 12.1\%$; 11 reports)^{26,39,42,50,55,58,59,61,84,97,101} and parental marriage (OR, 0.58; 95% CI, 0.51-0.66; $I^2 = 40.6\%$; 9 reports).^{19,20,35,36,59,77,78,94,97} All indicators of socioeconomic disadvantage were associated with higher risk, including lower youth education, with pooled ORs ranging from 1.25 to 5.24 (Figure 5; eTable 24 in Supplement 1).


Discussion

In this systematic review and meta-analysis on protective and risk factors for suicide mortality in children to young adults aged 24 years, we synthesized evidence from 68 observational cohort or case-control studies, including 90 reports. We found that the greatest odds of death by suicide were aspects of the individual's mental health (mental health disorders and recent contact with mental health services), underscoring the role of psychiatric vulnerability. At the socioecological level, suicide mortality was associated with maltreatment, out-of-home placement, justice-system involvement, lower education, and overall socioeconomic disadvantage. Although protective factors were rarely examined, the available evidence suggested that growing up in a stable home, such as living with both parents and parental marriage, was associated with lower suicide risk. Taken together, these findings suggest that death by suicide could occur at the interface of psychiatric vulnerability and social adversity. However, existing evidence remains fragmented and disproportionately focused on downstream risk markers, with limited attention to modifiable protective factors and upstream prevention targets.

Youths with a diagnosis of any mental disorder had almost 10-fold higher odds of death by suicide than youths without a diagnosis. Our findings are in line with a prior systematic review and meta-analysis by Too et al¹⁰⁸ that covered all ages. On the basis of administrative health records, they reported a pooled RR of 7.5 for all mental disorders combined. They also found particularly high risks for psychotic disorders (RR, 13.2) and mood disorders (RR, 12.3). Our results for mood disorders in youths were similar (OR, 11.32). Combined with our results in youths, findings suggest that mental health disorders are associated with substantially elevated suicide risk across the life course.

Nevertheless, effect sizes for some disorders were somewhat larger than those of Too et al¹⁰⁸ in our youth-focused review, particularly schizophrenia (OR, 22.23), possibly reflecting methodological differences. Furthermore, personality disorders in adult populations, particularly

Figure 5. Forest Plot of Pooled Associations of Socioecological Determinants With Youth Suicide Mortality (Meta-Analysis)



OR indicates odds ratio.

borderline personality disorder, have been associated with elevated suicide risk.^{108,109} In our review, borderline personality disorder was examined in only 1 psychological autopsy study in youths aged 18 to 25 years (OR, 12.66),¹⁹ highlighting a major evidence gap that may reflect ongoing controversy on diagnosing personality disorders during adolescence.¹¹⁰

A key clinical marker of elevated risk was prior self-harm. Although not a mental disorder in itself, self-harm was associated with death by suicide in youths (OR, 14.06). Other clinical markers in youths included recent contact with mental health care services, with especially high odds for the year prior to death (>7-fold elevated odds) rather than lifetime contacts (>3-fold elevated odds). We observed even higher odds of suicide mortality risk associated with psychiatric admission to hospital, but were unable to examine timing. Prior studies in adult populations have indicated that risk was particularly elevated in the period immediately following psychiatric discharge,¹¹¹ highlighting this transition as a critical window for prevention. Although our data cannot directly examine this period, this elevated risk may reflect a mismatch between ongoing vulnerability and the reduction or discontinuity of care following discharge. Evidence from our narrative synthesis suggests that service contact can occur across multiple settings. Future studies should therefore broaden their resources to include community-based and school-based contexts to better identify potentially missed prevention opportunities. Notably, service contact may reflect opportunities for intervention across the full continuum of care.

Of the 59 socioecological determinants vote-counted and 14 meta-analyzed, most reflected family-level socioeconomic status, such as parental education and household income. Heterogeneity was high, likely because of inconsistencies in exposure definitions across studies (eg, varying thresholds for low education). Socioeconomic disadvantage is a well-established risk factor for death by suicide in adulthood,^{7,112} and parental disadvantage is associated with suicidal distress in youths, including higher rates of emergency department visits for suicidal ideation and attempts.¹¹³ Studies clarifying the mechanisms whereby family-level and individual-level socioeconomic disadvantage is linked to death by suicide in youths could provide critical insights for prevention.

Adverse childhood experiences—such as maltreatment and bullying—are documented risk factors associated with suicidal distress in youths.³ Consistent with the literature, maltreatment and out-of-home placement emerged as the strongest socioecological determinants of death by suicide in our review. Evidence for bullying remained limited, however, and needs further elucidating. The juvenile justice system also arose as an essential element in targeted prevention. Youths involved in the system had an associated increased risk for suicide mortality. Adverse experiences tend to begin early in life and tend to recur, all while initiating developmental cascades that increase vulnerability, particularly with repeated exposure and disrupted access to protective factors. An in-depth understanding of these pathways is necessary to effect change and to identify points of preventive intervention to avoid long-term sequelae.

Protective factors were rarely examined in the studies we analyzed, consistent with findings from umbrella reviews of systematic reviews on social determinants of health.⁸ Of protective factors meta-analyzed, only 2 were consistently associated with lower suicide mortality: living with both parents and parental marriage. Although they are imperfect proxies of stability and social support, their consistent protective associations with suicide risk underscore the importance of strengthening family ties and relationship-based supports as part of youth suicide prevention efforts.

We noted a striking lack of research in populations known to be highly vulnerable, such as Indigenous youths, sexually and gender-diverse youths, and youths in low- and middle-income countries. Targeted epidemiological studies and prevention research are crucial for these underserved groups.

Our systematic review suggests that many of the risk and protective factors identified for suicide mortality may overlap with those reported for suicidal ideation and self-harm.⁵ The literature on death by suicide, however, appears more strongly focused on severe psychiatric disorders and less on socioecological determinants. This focus may reflect methodological differences, as studies of suicidal ideation and self-harm typically rely on self-reports, which are better suited to capturing

socioecological exposures such as maltreatment and bullying, whereas suicide mortality studies more frequently use psychological autopsies or register-based data, which provide more limited contextual information.

Avenues for suicide prevention among youths may concentrate on upstream strategies, including promoting stability in childhood environments and friendship building through engagement in schools and extracurricular activities; fostering positive attitudes toward learning and academic performance, and increased educational resources and support staff, for lifetime achievement; a reduction in childhood adversity and social disadvantage across the life course; and a rethinking of child protection such as out-of-home placement and youth justice involvement.¹¹⁰⁻¹¹³ Downstream strategies should focus on early detection, treatment, and continuity of care for special needs and mental health disorders, across all settings including school and health care services, and picking up on clinical cues such as self-harm.

Limitations and Strengths

Several limitations should be considered when interpreting these findings. First, we observed substantial heterogeneity (high I^2 values) among studies for most risk and protective factors, likely reflecting variations in design, time period, country, and exposure. Because of the small number of reports in each category, we could not conduct subgroup analyses (except by study design) to investigate sources of heterogeneity and interpretation of publication bias was limited. We were also unable to examine sex and gender differences, as most studies did not stratify by sex and gender. Second, the evidence originated largely from high-income and upper-middle-income countries, with very limited data from low-income and lower-middle-income settings. Similarly, vulnerable populations such as Indigenous or sexually and gender-diverse youths, were underrepresented. As a result, the identified risk and protective factors may not fully reflect context-specific determinants of suicide mortality in regions or communities with different social, cultural, and health care characteristics. Third, despite rigorous and systematic extraction of data into defined domains and factors, certain items may have been missed or incorrectly classified. Although all extracted data were verified by a second reviewer, extraction was not performed independently. Fourth, given the number and heterogeneity of factors, we relied on vote counting and minimally adjusted meta-analyses (age and sex). Findings are descriptive and do not capture independent associations or developmental pathways or interactions between factors. Fifth, we included only confirmed suicide deaths; deaths classified as other or as undetermined cause were excluded. In fact, some may have been suicides misclassified. Sixth, several factors associated with suicidal ideation and self-harm in young people, such as bullying,¹¹³ could not be meta-analyzed due to the paucity of available studies. Seventh, protective factors such as access to mental health care, faith and spirituality, and community belonging, were not documented.

The strengths of the present systematic review include the most comprehensive meta-analysis of current deaths by suicide in youth to date. Previously, most research focused on suicidal ideation or self-harm, or a combination thereof, but not strictly mortality. Many studies also examined various age ranges, but not strictly children to young adults younger than 25 years. With almost 100 high-quality reports from about 70 studies, we examined key risk and protective factors across all domains, from studies published in any language, worldwide.

Conclusions

In this systematic review and meta-analysis of risk and/or protective factors for suicide mortality in youth, the strongest risk factors that emerged were mental health disorders, contact with mental health care services in the previous year, maltreatment, out-of-home placement, and youth justice involvement. The strongest protective factor, or positive social determinant, was family stability. Further research is needed to clarify developmental pathways linking socioecological determinants to suicide mortality, including their interplay with mental health disorders and with bullying.

Research should also prioritize underrepresented Indigenous and sexually and gender-diverse populations for innovative preventive initiatives.

ARTICLE INFORMATION

Accepted for Publication: May 11, 2026.

Published: July 2, 2026. doi:10.1001/jamanetworkopen.2026.21622

Open Access: This is an open access article distributed under the terms of the [CC-BY-NC-ND License](#), which does not permit alteration or commercial use, including those for text and data mining, AI training, and similar technologies. © 2026 Geoffroy MC et al. *JAMA Network Open*.

Corresponding Author: Marie-Claude Geoffroy, PhD, Department of Psychiatry & Douglas Mental Health University Institute at McGill University, FBC, 6875 Blvd Lasalle, Montréal, QC H4H 1R3, Canada (marie-claude.geoffroy@mcgill.ca).

Author Affiliations: Douglas Mental Health University Institute, Department of Psychiatry, McGill University, Montreal, Quebec, Canada (Geoffroy, Nitu, Civita, Boruff, MacDonald, Anzarouth, Chartrand, Iyer, Thombs); Department of Epidemiology, Biostatistics, and Occupational Health, McGill University, Montreal, Quebec, Canada (Geoffroy, Baker-Sullivan, Thombs); The School of Physical and Occupational Therapy, McGill University, Montreal, Quebec, Canada (Boruff); The School of Communication Sciences and Disorders, McGill University, Montreal, Quebec, Canada (Boruff); Department of Psychiatry, Sunnybrook Health Science Centre, Toronto, Ontario, Canada (Sinyor); Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada (Sinyor); International Association for Suicide Prevention, Centre for Youth Mental Health (The Orygen Institute), Faculty of Medicine, Dentistry and Health Sciences, Melbourne University, Parkville, Victoria, Australia (Robinson); Division of Social and Transcultural Psychiatry, Department of Epidemiology, Biostatistics, and Occupational Health, McGill University, Montreal, Quebec, Canada (Iyer); Department of Educational and Counselling Psychology, McGill University, Montreal, Quebec, Canada (Langevin); The Centre for Mental Health, Melbourne School of Population and Global Health, Melbourne University, Parkville, Victoria, Australia (Pirkis); The Lady Davis Institute for Medical Research, Jewish General Hospital, Montreal, Quebec, Canada (Thombs).

Author Contributions: Dr Geoffroy and Ms Baker-Sullivan had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Geoffroy, Sinyor, Robinson, Iyer, Langevin, Pirkis.

Acquisition, analysis, or interpretation of data: Geoffroy, Baker-Sullivan, Nitu, Civita, Boruff, Sinyor, Robinson, MacDonald, Anzarouth, Chartrand, Iyer, Langevin, Thombs.

Drafting of the manuscript: Geoffroy, Boruff, Chartrand.

Critical review of the manuscript for important intellectual content: Geoffroy, Baker-Sullivan, Nitu, Civita, Boruff, Sinyor, Robinson, MacDonald, Anzarouth, Iyer, Langevin, Pirkis, Thombs.

Statistical analysis: Geoffroy, Baker-Sullivan.

Obtained funding: Geoffroy.

Administrative, technical, or material support: Geoffroy, Baker-Sullivan, Nitu, Civita, MacDonald, Anzarouth, Thombs.

Supervision: Geoffroy, Robinson.

Conflict of Interest Disclosures: Dr Sinyor reported receiving personal fees from Sunnybrook Health Sciences Centre and a Department of Psychiatry Academic Scholar Award for salary support during the conduct of the study. No other disclosures were reported.

Funding/Support: This work was supported by the Canada Research Chair in Youth Suicide Prevention and Canadian Institutes of Health Research (Dr Geoffroy).

Role of the Funder/Sponsor: The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Sharing Statement: See [Supplement 2](#).

Additional Contributions: We thank Danielle Buch (medical writer), for critical revision and substantive editing of the discussion and abstract. Compensation was provided for this work. Written permission to include this individual has been obtained.

REFERENCES

1. Bertuccio P, Amerio A, Grande E, et al. Global trends in youth suicide from 1990 to 2020: an analysis of data from the WHO mortality database. *EClinicalMedicine*. 2024;70:102506. doi:10.1016/j.eclinm.2024.102506
2. Chen PJ, Mackes N, Sacchi C, et al. Parental education and youth suicidal behaviours: a systematic review and meta-analysis. *Epidemiol Psychiatr Sci*. 2022;31:e19. doi:10.1017/S204579602200004X
3. Angelakis I, Austin JL, Gooding P. Association of childhood maltreatment with suicide behaviors among young people: a systematic review and meta-analysis. *JAMA Netw Open*. 2020;3(8):e2012563. doi:10.1001/jamanetworkopen.2020.12563
4. Santos B, Bettin BPC, de Almeida RMM. Determinants of risk for adolescent self-harm and suicide: a systematic review and diagram of shared and unique factors in non-clinical samples. *Neurosci Biobehav Rev*. 2025;177:106323. doi:10.1016/j.neubiorev.2025.106323
5. Richardson R, Connell T, Foster M, et al. Risk and protective factors of self-harm and suicidality in adolescents: an umbrella review with meta-analysis. *J Youth Adolesc*. 2024;53(6):1301-1322. doi:10.1007/s10964-024-01969-w
6. Favril L, Yu R, Uyar A, Sharpe M, Fazel S. Risk factors for suicide in adults: systematic review and meta-analysis of psychological autopsy studies. *Evid Based Ment Health*. 2022;25(4):148-155. doi:10.1136/ebmental-2022-300549
7. Favril L, Yu R, Geddes JR, Fazel S. Individual-level risk factors for suicide mortality in the general population: an umbrella review. *Lancet Public Health*. 2023;8(11):e868-e877. doi:10.1016/S2468-2667(23)00207-4
8. Na PJ, Shin J, Kwak HR, et al. Social determinants of health and suicide-related outcomes: a review of meta-analyses. *JAMA Psychiatry*. 2025;82(4):337-346. doi:10.1001/jamapsychiatry.2024.4241
9. Sawyer SM, Afifi RA, Bearinger LH, et al. Adolescence: a foundation for future health. *Lancet*. 2012;379(9826):1630-1640. doi:10.1016/S0140-6736(12)60072-5
10. Miranda-Mendizabal A, Castellví P, Parés-Badell O, et al. Gender differences in suicidal behavior in adolescents and young adults: systematic review and meta-analysis of longitudinal studies. *Int J Public Health*. 2019;64(2):265-283. doi:10.1007/s00038-018-1196-1
11. Boruff J. Search strategy for youth suicide mortality. McGill Librarian KS Psychiatry Search Sub-Repository. Posted online March 10, 2025. Accessed May 28, 2026. doi:10.5683/SP3/XCJCBT
12. Wells GA, Shea B, O'Connell D, et al. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomised Studies in Meta-Analyses*. Ottawa Hospital Research Institute; 2000.
13. American Psychiatric Association. *Report of the Presidential Task Force on the Social Determinants of Mental Health*. American Psychiatric Association; 2022.
14. Cochrane. Cochrane handbook for systematic reviews of interventions, version 6.5. 2024. Accessed January 16, 2026. <https://www.cochrane.org/authors/handbooks-and-manuals/handbook/current>
15. VanderWeele TJ. Optimal approximate conversions of odds ratios and hazard ratios to risk ratios. *Biometrics*. 2020;76(3):746-752. doi:10.1111/biom.13197
16. Cummings P. The relative merits of risk ratios and odds ratios. *Arch Pediatr Adolesc Med*. 2009;163(5):438-445. doi:10.1001/archpediatrics.2009.31
17. Lize SE, Scheyett AM, Morgan CR, Proescholdbell SK, Norwood T, Edwards D. Violent death rates and risk for released prisoners in North Carolina. *Violence Vict*. 2015;30(6):1019-1036. doi:10.1891/0886-6708.VV-D-13-00137
18. Weissman MM, Wolk S, Goldstein RB, et al. Depressed adolescents grown up. *JAMA*. 1999;281(18):1707-1713. doi:10.1001/jama.281.18.1707
19. Werbart Törnblom A, Sorjonen K, Runeson B, Rydelius PA. Who is at risk of dying young from suicide and sudden violent death? common and specific risk factors among children, adolescents, and young adults. *Suicide Life Threat Behav*. 2020;50(4):757-777. doi:10.1111/sltb.12614
20. Agerbo E, Nordentoft M, Mortensen PB. Familial, psychiatric, and socioeconomic risk factors for suicide in young people: nested case-control study. *BMJ*. 2002;325(7355):74. doi:10.1136/bmj.325.7355.74
21. Amin R, Mittendorfer-Rutz E, Mehlum L, et al. Does country of resettlement influence the risk of suicide in refugees? a case-control study in Sweden and Norway. *Epidemiol Psychiatr Sci*. 2021;30:e62. doi:10.1017/S2045796021000512
22. Beautrais AL. Suicide and serious suicide attempts in youth: a multiple-group comparison study. *Am J Psychiatry*. 2003;160(6):1093-1099. doi:10.1176/appi.ajp.160.6.1093

23. Beckman K, Mittendorfer-Rutz E, Lichtenstein P, et al. Mental illness and suicide after self-harm among young adults: long-term follow-up of self-harm patients, admitted to hospital care, in a national cohort. *Psychol Med*. 2016;46(16):3397-3405. doi:10.1017/S0033291716002282
24. Bjorkenstam C, Kosidou K, Bjorkenstam E. Childhood adversity and risk of suicide: cohort study of 548 721 adolescents and young adults in Sweden. *BMJ*. 2017;357:j1334. doi:10.1136/bmj.j1334
25. Blakely TA, Collings SCD, Atkinson J. Unemployment and suicide: evidence for a causal association? *J Epidemiol Community Health*. 2003;57(8):594-600. doi:10.1136/jech.57.8.594
26. Brent DA, Perper JA, Moritz G, et al. Psychiatric risk factors for adolescent suicide: a case-control study. *J Am Acad Child Adolesc Psychiatry*. 1993;32(3):521-529. doi:10.1097/00004583-199305000-00006
27. Brent DA, Perper JA, Moritz G, et al. Stressful life events, psychopathology, and adolescent suicide: a case control study. *Suicide Life Threat Behav*. 1993;23(3):179-187. doi:10.1111/j.1943-278X.1993.tb00178.x
28. Brent DA, Perper JA, Moritz G, Baugher M, Schweers J, Roth C. Firearms and adolescent suicide: a community case-control study. *Am J Dis Child*. 1993;147(10):1066-1071. doi:10.1001/archpedi.1993.02160340052013
29. Brent DA, Perper JA, Moritz G, et al. Familial risk factors for adolescent suicide: a case-control study. *Acta Psychiatr Scand*. 1994;89(1):52-58. doi:10.1111/j.1600-0447.1994.tb01485.x
30. Brent DA, Moritz G, Liotus L, et al. Familial risk factors for adolescent suicide: a case-control study. In: Kosky RJ, Eshkevari HS, Goldney RD, Hassan R, eds. *Suicide Prevention: The Global Context*. Springer; 1998:41-50.
31. Brent DA, Baugher M, Bridge J, Chen T, Chiappetta L. Age- and sex-related risk factors for adolescent suicide. *J Am Acad Child Adolesc Psychiatry*. 1999;38(12):1497-1505. doi:10.1097/00004583-199912000-00010
32. Burrell LV, Mehlum L, Qin P. Sudden parental death from external causes and risk of suicide in the bereaved offspring: a national study. *J Psychiatr Res*. 2018;96:49-56. doi:10.1016/j.jpsychires.2017.09.023
33. Chartier MJ, Brownell M, Star L, et al. The mental health of First Nations children in Manitoba: a population-based retrospective cohort study using linked administrative data. *Can J Psychiatry*. 2024;69(6):404-414. doi:10.1177/07067437241226998
34. Chen EYH, Chan WSC, Wong PWC, et al. Suicide in Hong Kong: a case-control psychological autopsy study. *Psychol Med*. 2006;36(6):815-825. doi:10.1017/S0033291706007240
35. Chen YY, Gunnell D, Lu CL, Chang SS, Lu TH, Li CY. Perinatal risk factors for suicide in young adults in Taiwan. *Int J Epidemiol*. 2013;42(5):1381-1389. doi:10.1093/ije/dyt129
36. Cheng CCJ, Yen WJ, Chang WT, Wu KCC, Ko MC, Li CY. Risk of adolescent offspring's completed suicide increases with prior history of their same-sex parents' death by suicide. *Psychol Med*. 2014;44(9):1845-1854. doi:10.1017/S0033291713002298
37. Cheng CM, Chang WH, Tsai SJ, et al. Risk of all-cause and suicide death in patients with schizophrenia: an entire-population longitudinal study in Taiwan. *J Clin Psychiatry*. 2023;84(6):22m14747. doi:10.4088/JCP.22m14747
38. Clapperton A, Newstead S, Bugeja L, Pirkis J. Relative risk of suicide following exposure to recent stressors, Victoria, Australia. *Aust N Z J Public Health*. 2019;43(3):254-260. doi:10.1111/1753-6405.12886
39. Cnattingius S, Svensson T, Granath F, Iliadou A. Maternal smoking during pregnancy and risks of suicidal acts in young offspring. *Eur J Epidemiol*. 2011;26(6):485-492. doi:10.1007/s10654-011-9556-7
40. Cumming C, Bell MF, Segal L, et al. Maternal incarceration increases the risk of self-harm but not suicide: a matched cohort study. *Epidemiol Psychiatr Sci*. 2023;32:e33. doi:10.1017/S2045796023000264
41. Cybulski L, Ashcroft DM, Carr MJ, et al. Risk factors for nonfatal self-harm and suicide among adolescents: two nested case-control studies conducted in the UK Clinical Practice Research Datalink. *J Child Psychol Psychiatry*. 2022;63(9):1078-1088. doi:10.1111/jcpp.13552
42. Davidson LE, Rosenberg ML, Mercy JA, Franklin J, Simmons JT. An epidemiologic study of risk factors in two teenage suicide clusters. *JAMA*. 1989;262(19):2687-2692. doi:10.1001/jama.1989.03430190071034
43. Del Pozo Banos M, Lloyd K, Dennis M, Gunnell D, Scourfield J, John A. Case-control study of suicide in children and young people using linked primary and secondary routinely collected electronic health records. *Eur Psychiatry*. 2018;48(suppl 1):S117. doi:10.1016/j.eurpsy.2017.12.022
44. Drucker AM, Thiruchelvam D, Redelmeier DA. Eczema and subsequent suicide: a matched case-control study. *BMJ Open*. 2018;8(11):e023776. doi:10.1136/bmjopen-2018-023776
45. Edwards AC, Ohlsson H, Lannoy S, et al. Exposure to alcohol outlets and risk of suicidal behavior in a Swedish cohort of young adults. *Alcohol Clin Exp Res (Hoboken)*. 2023;47(5):930-939. doi:10.1111/acer.15051

46. Evensen M, Klitkou ST, Tollånes MC, Júlíusson PB, Kravdal Ø. Parental income gradients in child and adolescent mortality: Norwegian trends over half a century. *Scand J Public Health*. 2024;52(4):402-409. doi:10.1177/14034948231151990
47. Fazel S, Wolf A, Långström N, Newton CR, Lichtenstein P. Premature mortality in epilepsy and the role of psychiatric comorbidity: a total population study. *Lancet*. 2013;382(9905):1646-1654. doi:10.1016/S0140-6736(13)60899-5
48. Fazel S, Wolf A, Pillas D, Lichtenstein P, Långström N. Suicide, fatal injuries, and other causes of premature mortality in patients with traumatic brain injury: a 41-year Swedish population study. *JAMA Psychiatry*. 2014;71(3):326-333. doi:10.1001/jamapsychiatry.2013.3935
49. Feigelman W, Joiner T, Rosen Z, Silva C, Mueller AS. Contrasts between young males dying by suicide, those dying from other causes and those still living: observations from the national longitudinal survey of adolescent to adult health. *Arch Suicide Res*. 2016;20(3):389-401. doi:10.1080/13811118.2015.1104270
50. Feigelman W, Joiner T, Rosen Z, Silva C. Investigating correlates of suicide among male youth: questioning the close affinity between suicide attempts and deaths. *Suicide Life Threat Behav*. 2016;46(2):191-205. doi:10.1111/sltb.12183
51. Feigelman W, Plöderl M, Rosen Z, Cerel J. Research note on whether sexual minority individuals are over-represented among suicide's casualties. *Crisis*. 2020;41(3):229-232. doi:10.1027/0227-5910/a000626
52. Feigelman W, Rosen Z, Cerel J. Unraveling the complex web of associations between easy access to firearms and premature mortalities. *Suicide Life Threat Behav*. 2020;50(1):277-291. doi:10.1111/sltb.12580
53. Feigelman W, Coleman D, Rosen Z. Examining the social origins and young adult life trajectories of high traditional masculinity (HTM) males: a group at elevated suicide risk. *Suicide Life Threat Behav*. 2021;51(4):696-714. doi:10.1111/sltb.12753
54. Fitzgerald C, Dalsgaard S, Nordentoft M, Erlangsen A. Suicidal behaviour among persons with attention-deficit hyperactivity disorder. *Br J Psychiatry*. 2019;215(4):615-620. doi:10.1192/bjp.2019.128
55. Freuchen A, Kjelsberg E, Lundervold AJ, Grøholt B. Differences between children and adolescents who commit suicide and their peers: a psychological autopsy of suicide victims compared to accident victims and a community sample. *Child Adolesc Psychiatry Ment Health*. 2012;6:1. doi:10.1186/1753-2000-6-1
56. Goldstein TR, Bridge JA, Brent DA. Sleep disturbance preceding completed suicide in adolescents. *J Consult Clin Psychol*. 2008;76(1):84-91. doi:10.1037/0022-006X.76.1.84
57. Gould MS, Shaffer D, Davies M. Truncated pathways from childhood to adulthood: attrition in follow-up studies due to death. In: Robins LN, Rutter M, eds. *Straight and Devious Pathways From Childhood to Adulthood*. Cambridge University Press; 1990:3-9.
58. Gould MS, Fisher P, Parides M, Flory M, Shaffer D. Psychosocial risk factors of child and adolescent completed suicide. *Arch Gen Psychiatry*. 1996;53(12):1155-1162. doi:10.1001/archpsyc.1996.01830120095016
59. Grande E, Vichi M, Alicandro G, et al. Suicide among adolescents in Italy: a nationwide cohort study of the role of family characteristics. *Eur Child Adolesc Psychiatry*. 2021;30(7):1037-1045. doi:10.1007/s00787-020-01591-8
60. Gray D, Achilles J, Keller T, et al. Utah youth suicide study, phase I: government agency contact before death. *J Am Acad Child Adolesc Psychiatry*. 2002;41(4):427-434. doi:10.1097/00004583-200204000-00015
61. Grøholt B, Ekeberg O, Wichstrøm L, Haldorsen T. Youth suicide in Norway, 1990-1992: a comparison between children and adolescents completing suicide and age- and gender-matched controls. *Suicide Life Threat Behav*. 1997;27(3):250-263. doi:10.1111/j.1943-278X.1997.tb00407.x
62. Hjern A, Lindblad F, Vinnerljung B. Suicide, psychiatric illness, and social maladjustment in intercountry adoptees in Sweden: a cohort study. *Lancet*. 2002;360(9331):443-448. doi:10.1016/S0140-6736(02)09674-5
63. Huang WC, Hsu CY, Chang CM, et al. Psychiatrist density and risk of suicide: a multilevel case-control study based on a national sample in Taiwan. *Psychiatry Clin Neurosci*. 2024;78(1):69-76. doi:10.1111/pcn.13607
64. Hung YA, Liao SC, Chang CM, et al. Population-attributable risk of psychiatric disorders for suicide among adolescents and young adults in Taiwan. *Psychol Med*. 2023;53(13):6161-6170. doi:10.1017/S0033291722003361
65. Jokiranta-Olkoniemi E, Gyllenberg D, Sucksdorff D, et al. Risk for premature mortality and intentional self-harm in autism spectrum disorders. *J Autism Dev Disord*. 2021;51(9):3098-3108. doi:10.1007/s10803-020-04768-x
66. Jonsson U, Alexanderson K, Kjeldgård L, Mittendorfer-Rutz E. Psychiatric diagnoses and risk of suicidal behaviour in young disability pensioners: prospective cohort studies of all 19-23 year olds in Sweden in 1995, 2000, and 2005, respectively. *PLoS One*. 2014;9(11):e111618. doi:10.1371/journal.pone.0111618

67. Kaplan MS, Huguet N, McFarland BH, et al. Use of alcohol before suicide in the United States. *Ann Epidemiol*. 2014;24(8):588-592. doi:10.1016/j.annepidem.2014.05.008
68. Knipe DW, Gunnell D, Pieris R, et al. Socioeconomic position and suicidal behaviour in rural Sri Lanka: a prospective cohort study of 168,000+ people. *Soc Psychiatry Psychiatr Epidemiol*. 2019;54(7):843-855. doi:10.1007/s00127-019-01672-3
69. Knipe D, Moran P, Howe LD, et al. Ethnicity and suicide in England and Wales: a national linked cohort study. *Lancet Psychiatry*. 2024;11(8):611-619. doi:10.1016/S2215-0366(24)00184-6
70. Kong Y, Zhang J. Access to farming pesticides and risk for suicide in Chinese rural young people. *Psychiatry Res*. 2010;179(2):217-221. doi:10.1016/j.psychres.2009.12.005
71. Kwan YK, Ip WC. Suicidality and migration among adolescents in Hong Kong. *Death Stud*. 2007;31(1):45-66. doi:10.1080/07481180600993144
72. Lee WC, Lai JY, Pan CH, et al. Healthcare utilization, psychiatric disorders, and physical illnesses shortly before suicide mortality in adolescents in Taiwan. *Psychol Med*. 2023;53(7):2885-2894. doi:10.1017/S0033291721004864
73. Li CI, Daling JR, Emanuel I. Birthweight and risk of overall and cause-specific childhood mortality. *Paediatr Perinat Epidemiol*. 2003;17(2):164-170. doi:10.1046/j.1365-3016.2003.00487.x
74. Liu HL, Chen LH, Huang SM. Outpatient health care utilization of suicide decedents in their last year of life. *Suicide Life Threat Behav*. 2012;42(4):445-452. doi:10.1111/j.1943-278X.2012.00103.x
75. Mittendorfer-Rutz E, Rasmussen F, Wasserman D. Restricted fetal growth and adverse maternal psychosocial and socioeconomic conditions as risk factors for suicidal behaviour of offspring: a cohort study. *Lancet*. 2004;364(9440):1135-1140. doi:10.1016/S0140-6736(04)17099-2
76. Na EJ, Lee H, Myung W, et al. Risks of completed suicide of community individuals with ICD-10 disorders across age groups: a nationwide population-based nested case-control study in South Korea. *Psychiatry Investig*. 2019;16(4):314-324. doi:10.30773/pi.2019.02.19
77. Neugebauer R, Reuss ML. Association of maternal, antenatal and perinatal complications with suicide in adolescence and young adulthood. *Acta Psychiatr Scand*. 1998;97(6):412-418. doi:10.1111/j.1600-0447.1998.tb10024.x
78. Ortin-Peralta A, Keski-Säntti M, Gissler M, Veijola J, Sourander A, Duarte CS. Parental suicide attempts and offspring's risk of attempting or dying by suicide: does the timing of a parental suicide attempt matter? *Psychol Med*. 2023;53(3):977-986. doi:10.1017/S0033291721002397
79. Palmer L, Prindle J, Putnam-Hornstein E. A population-based examination of suicide and child protection system involvement. *J Adolesc Health*. 2021;69(3):465-469. doi:10.1016/j.jadohealth.2021.02.006
80. Park JY, Moon KT, Chae YM, Jung SH. Effect of sociodemographic factors, cancer, psychiatric disorder on suicide: gender and age-specific patterns. *J Prev Med Public Health*. 2008;41(1):51-60. doi:10.3961/jpmph.2008.41.1.51
81. Qin P, Mortensen PB, Pedersen CB. Frequent change of residence and risk of attempted and completed suicide among children and adolescents. *Arch Gen Psychiatry*. 2009;66(6):628-632. doi:10.1001/archgenpsychiatry.2009.20
82. Qin P. Suicide risk in relation to level of urbanicity: a population-based linkage study. *Int J Epidemiol*. 2005;34(4):846-852. doi:10.1093/ije/dyi085
83. Renaud J, Berlim MT, McGirr A, Tousignant M, Turecki G. Current psychiatric morbidity, aggression/impulsivity, and personality dimensions in child and adolescent suicide: a case-control study. *J Affect Disord*. 2008;105(1-3):221-228. doi:10.1016/j.jad.2007.05.013
84. Renaud J, Berlim MT, Séguin M, McGirr A, Tousignant M, Turecki G. Recent and lifetime utilization of health care services by children and adolescent suicide victims: a case-control study. *J Affect Disord*. 2009;117(3):168-173. doi:10.1016/j.jad.2009.01.004
85. Renaud J, Berlim MT, Begolli M, McGirr A, Turecki G. Sexual orientation and gender identity in youth suicide victims: an exploratory study. *Can J Psychiatry*. 2010;55(1):29-34. doi:10.1177/070674371005500105
86. Rhodes AE, Boyle MH, Bridge JA, et al. The medical care of male and female youth who die by suicide: a population-based case control study. *Can J Psychiatry*. 2018;63(3):161-169. doi:10.1177/0706743717741060
87. Rojas Y, Stenberg SA. Early life circumstances and male suicide—a 30-year follow-up of a Stockholm cohort born in 1953. *Soc Sci Med*. 2010;70(3):420-427. doi:10.1016/j.socscimed.2009.10.026

88. Saunders NR, Lebenbaum M, Stukel TA, et al. Suicide and self-harm trends in recent immigrant youth in Ontario, 1996-2012: a population-based longitudinal cohort study. *BMJ Open*. 2017;7(9):e014863. doi:[10.1136/bmjopen-2016-014863](https://doi.org/10.1136/bmjopen-2016-014863)
89. Seguin M, Renaud J, Lesage A, Robert M, Turecki G. Youth and young adult suicide: a study of life trajectory. *J Psychiatr Res*. 2011;45(7):863-870. doi:[10.1016/j.jpsychires.2011.05.005](https://doi.org/10.1016/j.jpsychires.2011.05.005)
90. Shaffer D, Fisher P, Hicks RH, Parides M, Gould M. Sexual orientation in adolescents who commit suicide. *Suicide Life Threat Behav*. 1995;25(S1):64-71. doi:[10.1111/j.1943-278X.1995.tb00491.x](https://doi.org/10.1111/j.1943-278X.1995.tb00491.x)
91. Shaffer D, Gould MS, Fisher P, et al. Psychiatric diagnosis in child and adolescent suicide. *Arch Gen Psychiatry*. 1996;53(4):339-348. doi:[10.1001/archpsyc.1996.01830040075012](https://doi.org/10.1001/archpsyc.1996.01830040075012)
92. Shafii M, Carrigan S, Whittinghill JR, Derrick A. Psychological autopsy of completed suicide in children and adolescents. *Am J Psychiatry*. 1985;142(9):1061-1064. doi:[10.1176/ajp.142.9.1061](https://doi.org/10.1176/ajp.142.9.1061)
93. Shafii M, Steltz-Lenarsky J, Derrick AM, Beckner C, Whittinghill JR. Comorbidity of mental disorders in the post-mortem diagnosis of completed suicide in children and adolescents. *J Affect Disord*. 1988;15(3):227-233. doi:[10.1016/0165-0327\(88\)90020-1](https://doi.org/10.1016/0165-0327(88)90020-1)
94. Shah S, Hoffman RE, Wake L, Marine WM. Adolescent suicide and household access to firearms in Colorado: results of a case-control study. *J Adolesc Health*. 2000;26(3):157-163. doi:[10.1016/S1054-139X\(99\)00064-6](https://doi.org/10.1016/S1054-139X(99)00064-6)
95. Sohn J, Cho J, Moon KT, et al. Medical care expenditure in suicides from non-illness-related causes. *J Prev Med Public Health*. 2014;47(6):327-335. doi:[10.3961/jpmph.14.038](https://doi.org/10.3961/jpmph.14.038)
96. Sörberg Wallin A, Sjöqvist H, Dehara M, et al. Parenthood and lower risk of suicide in women and men: the total Swedish population followed across adulthood. *Soc Psychiatry Psychiatr Epidemiol*. 2022;57(11):2241-2250. doi:[10.1007/s00127-022-02321-y](https://doi.org/10.1007/s00127-022-02321-y)
97. Steck N, Egger M, Schimmelmann BG, Kupferschmid S; Swiss National Cohort. Suicide in adolescents: findings from the Swiss National cohort. *Eur Child Adolesc Psychiatry*. 2018;27(1):47-56. doi:[10.1007/s00787-017-1019-6](https://doi.org/10.1007/s00787-017-1019-6)
98. Tidemalm D, Beckman K, Dahlin M, et al. Age-specific suicide mortality following non-fatal self-harm: national cohort study in Sweden. *Psychol Med*. 2015;45(8):1699-1707. doi:[10.1017/S0033291714002827](https://doi.org/10.1017/S0033291714002827)
99. Webb RT, Pickles AR, Appleby L, Mortensen PB, Abel KM. Death by unnatural causes during childhood and early adulthood in offspring of psychiatric inpatients. *Arch Gen Psychiatry*. 2007;64(3):345-352. doi:[10.1001/archpsyc.64.3.345](https://doi.org/10.1001/archpsyc.64.3.345)
100. Webb RT, Qin P, Stevens H, Mortensen PB, Appleby L, Shaw J. National study of suicide in all people with a criminal justice history. *Arch Gen Psychiatry*. 2011;68(6):591-599. doi:[10.1001/archgenpsychiatry.2011.7](https://doi.org/10.1001/archgenpsychiatry.2011.7)
101. Weitoft GR, Hjern A, Haglund B, Rosén M. Mortality, severe morbidity, and injury in children living with single parents in Sweden: a population-based study. *Lancet*. 2003;361(9354):289-295. doi:[10.1016/S0140-6736\(03\)12324-0](https://doi.org/10.1016/S0140-6736(03)12324-0)
102. Werbart Törnblom A, Sorjonen K, Runeson B, Rydelius PA. Life events and coping strategies among young people who died by suicide or sudden violent death. *Front Psychiatry*. 2021;12:670246. doi:[10.3389/fpsy.2021.670246](https://doi.org/10.3389/fpsy.2021.670246)
103. Wyder M, Ward P, De Leo D. Separation as a suicide risk factor. *J Affect Disord*. 2009;116(3):208-213. doi:[10.1016/j.jad.2008.11.007](https://doi.org/10.1016/j.jad.2008.11.007)
104. Yao S, Kuja-Halkola R, Thornton LM, et al. Familial liability for eating disorders and suicide attempts: evidence from a population registry in Sweden. *JAMA Psychiatry*. 2016;73(3):284-291. doi:[10.1001/jamapsychiatry.2015.2737](https://doi.org/10.1001/jamapsychiatry.2015.2737)
105. Zambon F, Laflamme L, Spolaore P, Visentin C, Hasselberg M. Youth suicide: an insight into previous hospitalisation for injury and sociodemographic conditions from a nationwide cohort study. *Inj Prev*. 2011;17(3):176-181. doi:[10.1136/ip.2010.030080](https://doi.org/10.1136/ip.2010.030080)
106. Zhao S, Zhang J. Suicide risks among adolescents and young adults in rural China. *Int J Environ Res Public Health*. 2014;12(1):131-145. doi:[10.3390/ijerph120100131](https://doi.org/10.3390/ijerph120100131)
107. Ioannidis JPA, Trikalinos TA. The appropriateness of asymmetry tests for publication bias in meta-analyses: a large survey. *CMAJ*. 2007;176(8):1091-1096. doi:[10.1503/cmaj.060410](https://doi.org/10.1503/cmaj.060410)
108. Too LS, Spittal MJ, Bugeja L, Reifels L, Butterworth P, Pirkis J. The association between mental disorders and suicide: a systematic review and meta-analysis of record linkage studies. *J Affect Disord*. 2019;259:302-313. doi:[10.1016/j.jad.2019.08.054](https://doi.org/10.1016/j.jad.2019.08.054)
109. Lak M, Shakiba S, Dolatshahi B, Saatchi M, Shahrabaf M, Jafarpour A. The prevalence of suicide ideation, suicide attempt and suicide in borderline personality disorder patients: a systematic review and meta-analysis. *Gen Hosp Psychiatry*. 2025;95:52-61. doi:[10.1016/j.genhosppsy.2025.04.005](https://doi.org/10.1016/j.genhosppsy.2025.04.005)

110. Kaess M, Brunner R, Chanan A. Borderline personality disorder in adolescence. *Pediatrics*. 2014;134(4):782-793. doi:10.1542/peds.2013-3677
111. Aaltonen K, Sund R, Hakulinen C, Pirkola S, Isometsä E. Variations in suicide risk and risk factors after hospitalization for depression in Finland, 1996-2017. *JAMA Psychiatry*. 2024;81(5):506-515. doi:10.1001/jamapsychiatry.2023.5512
112. Balaj M, Henson CA, Aronsson A, et al; IHME-CHAIN Collaborators. Effects of education on adult mortality: a global systematic review and meta-analysis. *Lancet Public Health*. 2024;9(3):e155-e165. doi:10.1016/S2468-2667(23)00306-7
113. Holt MK, Vivolo-Kantor AM, Polanin JR, et al. Bullying and suicidal ideation and behaviors: a meta-analysis. *Pediatrics*. 2015;135(2):e496-e509. doi:10.1542/peds.2014-1864

SUPPLEMENT 1.

- eTable 1. Mental Health Disorders Included or Excluded for Vote-Counting
- eTable 2. Physical Health Disorders Included or Excluded for Vote-Counting
- eTable 3. Health-Care Service Contacts Included or Excluded for Vote-Counting
- eTable 4. Prenatal and Perinatal Factors Included or Excluded for Vote-Counting
- eTable 5: Socioecological Determinants Included or Excluded for Vote-Counting
- eTable 6. Original Assessment of Youth Educational Attainment and Harmonization
- eTable 7. Original Assessment of Maternal Age at Child's Birth and Harmonization
- eTable 8. Original Assessment of Maternal Educational Attainment and Harmonization
- eTable 9. Original Assessment of Paternal Educational Attainment and Harmonization
- eTable 10. Original Assessment of Parental Income and Harmonization
- eTable 11. Original Assessment of Residential Moves and Harmonization
- eTable 12. Original Assessment of Rurality and Harmonization
- eTable 13. Original Assessment of Living Arrangement and Harmonization
- eTable 14. Original Assessment of Parental Marital Status and Harmonization
- eTable 15. Original Assessment of Criminal Justice Involvement (Youth) and Harmonization
- eTable 16. Characteristics of Included Studies After Full-Text Screening, with Studies Included in the Meta-Analysis Shown in Bold
- eTable 17: Number of Reports Reporting Harmful, Protective, Null, or Unreported Associations Between Risk and/or Protective Factors and Youth Suicide Mortality (Vote-Counting Synthesis)
- eTable 18. Quality Ratings for Studies Included in the Meta-Analysis Using the Newcastle Ottawa Scale (Cohort Studies)
- eTable 19. Quality Ratings for Studies Included in the Meta-Analysis Using the Newcastle Ottawa Scale (Case-Control Studies)
- eTable 20. Sensitivity Subgroup Analyses by Study Design (Case-control vs Cohort) for Associations with Youth Suicide Mortality, Restricted to Subgroups with ≥ 2 Studies
- eTable 21. Study-Level Odds Ratios and 95% Confidence Intervals and Pooled Estimates for Mental Health Disorders and Suicide Mortality in Youth
- eTable 22. Study-Level Odds Ratios and 95% Confidence Intervals and Pooled Estimates for Physical Disorders and Suicide Mortality in Youth
- eTable 23. Study-Level Odds Ratios and 95% Confidence Intervals and Pooled Estimates for Health-Care Service Contacts and Suicide Mortality in Youth
- eTable 24. Study-Level Odds Ratios and 95% Confidence Intervals and Pooled Estimates for Socioecological Determinants and Suicide Mortality in Youth
- eFigure 1. Number of Reports Reporting Harmful, Protective, Null, or Unreported Associations Between Health Related Risk Factors and Youth Suicide Mortality (Vote-Counting Synthesis)
- eFigure 2. Funnel Plots for Individual Meta-Analyses of Risk and Protective Factors for Youth Suicide Mortality
- eMethods.
- eAppendix 1. Title and Abstract Screening Guide
- eAppendix 2. Full Text Screening Guide
- eAppendix 3. Newcastle-Ottawa Quality Assessment Form for Cohort Studies
- eAppendix 4. Newcastle-Ottawa Quality Assessment Form for Case-Control Studies
- eAppendix 5. Full List of Excluded Studies After Full-Text Screening

SUPPLEMENT 2.

Data Sharing Statement